

Vaccines for sheep and goats

Susan Schoenian, Sheep & Goat Specialist, University of Maryland Extension [last updated 4.19.2012]

Vaccine	Manufacturer	Approved Species	Specs	Sheep/Goat Dosage	Slaughter Withdrawal
Bar-Vac® CD-T	Boehringer	Cattle Sheep Goats	Vaccine for immunizing sheep, goats, and cattle against tetanus and overeating disease	2 ml SQ Repeat in 21 to 28 days Repeat annually	21 days
C & D Antitoxin	Boehringer- Ingleheim	Cattle Calves Sheep Suckling lambs Goats Baby pigs	Vaccine for prevention (lasting 3 weeks) and treatment of enterotoxemia	Sheep: 10 ml IM or SQ Suckling lambs: 3 ml For treatment, double above doses	21 days
C & D Toxoid	Colorado Serum	Cattle Sheep	For vaccination against enterotoxemia caused by the toxins of Clostridium perfringens type C and D.	2 ml SQ Repeat in 21 to 28 days Repeat annually	21 days
Case-Bac™	Colorado Serum	Sheep Goats	Vaccine to aid in the prevention and control of Caseous lymphadenitis	2 ml SQ (auxillary space) Repeat full 2 ml dose in 4 weeks. Annual booster	21 days
Campylobacter fetus bacterin	Colorado Serum	Sheep	Protect ewes against vibrio abortion	5 ml SQ on side of neck or rib cage shortly before exposure to ram, 2 nd dose 60 to 90 days later, repeat annually	21 days
Caseous D-T	Colorado Serum	Sheep	Vaccine against Caseous Lymphadenitis, overeating disease, and lock jaw	2 ml SQ to sheep of any age, repeat in 4 weeks, annual booster recommended	21 days

Vaccine	Manufacturer	Approved Species	Specs	Sheep/Goat Dosage	Slaughter Withdrawal
Chlamydia Psittaci Bacterin	Colorado Serum	Sheep Goats	Protect ewes against ovine enzootic abortion	2 ml SQ in the top of the neck 60 days prior to breeding, repeat 2 ml dose in 30 days, revaccinate annually just prior to breeding	60 days
Clostratox BCD	Novartis	Cattle Sheep	Aid in the prevention of Enterotoxemia caused by Clostridium perfringens Types C and D. The combination of these fractions may protect against Type B	Feeder lambs: 10 ml SQ Double dosage for treatment	21 days
Clostridium perfringens type C & D antitoxin	Colorado Serum	Suckling calves Feeder calves Suckling pigs Suckling lambs Suckling goats	Multivalent antitoxin to treat and prevent enterotoxemia caused by types B, C, and D	Suckling lambs and kids: 5 ml Suckling feeder lambs: 10 ml	21 days
Covexin®-8	Schering-Plough	Cattle Sheep	For the vaccination of healthy cattle and sheep against diseases caused by <i>Clostridium chauvoei</i> , <i>Cl. septicum</i> , <i>Cl. novyi</i> Type B, <i>Cl. haemolyticum</i> (known elsewhere as <i>Cl. novyi</i> Type D), <i>Cl. tetani</i> and <i>Cl. perfringens</i> Types C and D.	5 ml SQ, followed by 2 ml dose in six weeks. Repeat annually with 2 ml dose.	21 days
Ovine Ecthyma vaccine Live virus.	Colorado Serum	Sheep Goats	Protect healthy sheep and goats against soremouth	Apply to wool-free, scarified area Do not use within 24 hours of dipping or spraying	21 days
Essential 3 + T	Colorado Serum	Cattle Swine Sheep Goats	Aid in prevention of enterotoxemia caused by Cl. Perfringens Type B, C, and D and tetanus	2 ml SQ Repeat in 3-4 weeks Repeat annually	21 days

Vaccine	Manufacturer	Approved Species	Specs	Sheep/Goat Dosage	Slaughter Withdrawal
Pasteurella (pneumonia)	Colorado Serum	Cattle Sheep Goats	Aid in prevention of pneumonia caused by Mannheimia haemolytica and Pasteurella multocida	2 ml SQ Repeat in 2-4 weeks	21 days
Rabies vaccine Imrab 3 Imrab Large Animal	Merial	Cattle Sheep Horses Dogs Cats	Killed virus recommended for vaccination for prevention of disease due to rabies.	2 ml SQ Repeat in 1 year and every 3 years thereafter	21 days
Tetanus Antitoxin	Colorado Serum	Horses Cattle Sheep Swine Goats	For use whenever a non-immunized animal, or one whose immune status is unknown, suffers a deep penetrating wound that has or may become contaminated with soil. It provides quick but short-term protection.	Administer 1500 units (5 ml) SQ or IM for 7-14 days of passive immunity For treatment, administer 3000-15000 units. Administer 2 nd dose in 7 days	21 days
Tetanus Toxoid	Colorado Serum	Sheep Goats Swine Cattle Horses	Vaccine for establishing immunity against tetanus	2 doses of 1 ml per 100 lbs. IM each 30 days apart	21 days
Ultrabac 8	Pfizer Animal Health	Cattle Sheep	Aid in preventing disease caused by Cl. chauvoei, septicum, haemolyticum, novyi, sordellii and perfringens types C & D. Immunity may be provided against type B.	2.5 ml SQ Repeat in 4-6 weeks Repeat annually	21 days
Ultrabac CD	Pfizer Animal Health	Cattle Sheep	For the immunization of healthy cattle and sheep against Enterotoxemia Enteritis caused by Cl. perfringens types C & D.	1 ml IM or SQ Repeat in 4-6 weeks Repeat annually	21 days

Vaccine	Manufacturer	Approved Species	Specs	Sheep/Goat Dosage	Slaughter Withdrawal
UltraChoice 8	Pfizer Animal Health	Cattle Sheep	Aid in preventing diseases caused by Cl. chauvoei, septicum, novyi, sordellii, haemolyticum and perfringens types C and D	1 ml SQ in neck Repeat in 4-6 weeks Repeat annually	21 days
Vision® CD-T with Spur®	Intervet	Cattle Sheep Goats	Aid in the prevention of enterotoxemia caused by Clostridium perfringens types C & D	2 ml SQ Repeat dose in 3 to 4 weeks	21 days

Source: product labels